

Esther G.C. Troost, MD, PhD**Aswin L. Hoffmann, MSc**

Department of Radiation Oncology

(MAASTRO)

GROW School for Oncology and

Developmental Biology

Maastricht University Medical Centre

Maastricht, The Netherlands

Johan Bussink, MD, PhD

Department of Radiation Oncology

Radboud University Nijmegen

Medical Centre

Nijmegen, The Netherlands

REFERENCES

1. Vinogradskiy Y, Tucker SL, Bluett JB, Wages CA, Liao Z, Martel MK. Prescribing radiation dose to lung cancer patients based on personalized toxicity estimates. *J Thorac Oncol* 2012;7:1676–1682.
2. Govaert SL, Troost EG, Schuurbijs OC, et al. Treatment outcome and toxicity of intensity-modulated (chemo) radiotherapy in stage III non-small cell lung cancer patients. *Radiat Oncol* 2012;7:150.
3. Hoffmann AL, Troost EG, Huizenga H, Kaanders JH, Bussink J. Individualized dose prescription for hypofractionation in advanced non-small-cell lung cancer radiotherapy: an in silico trial. *Int J Radiat Oncol Biol Phys* 2012;83:1596–1602.
4. Partridge M, Ramos M, Sardaro A, Brada M. Dose escalation for non-small cell lung cancer: analysis and modelling of published literature. *Radiother Oncol* 2011;99:6–11.
5. van Baardwijk A, Wanders S, Boersma L, et al. Mature results of an individualized radiation dose prescription study based on normal tissue constraints in stages I to III non-small-cell lung cancer. *J Clin Oncol* 2010;28:1380–1386.

Two Rare Exon 21 EGFR Mutations in Patients Treated with Gefitinib

To the Editor:

Treatment of non-small-cell lung cancer patients with reversible inhibitors

Disclosure: The authors declare no conflict of interest.

Address for correspondence: Ederne Arriola, MD, PhD, Oncology Department, Hospital del Mar- Parc de Salut Mar, Passeig Marítim 25–29, 08003, Barcelona. E-mail: earriola@parcadesalutmar.cat

Copyright © 2013 by the International Association for the Study of Lung Cancer
ISSN: 1556-0864/13/0804-e36




FIGURE 1. Sequencing chromatograms showing EGFR mutations affecting cases 1 and 2. A, Case 1-arginine(R) to cysteine (C) substitution at amino acid position 836 (p.R836C) resulting from a CGC→TGC exchange. B, Case 2-threonine (T) to serine (S) substitution at amino acid position 854 (p.T854S) resulting from a ACA→TCA exchange.

of the tyrosine kinase domain of epidermal growth factor receptor (EGFR), gefitinib and erlotinib, in patients with EGFR mutations result in 70% to 80% responses.^{1,2} Ninety percent of these consist of deletions of exon 19 and p.L858R mutations on exon 21.² However, through diverse mechanisms³ resistance will eventually appear. There are also a minority of infrequent EGFR mutations at diagnosis whose predictive role is poorly characterized.⁴

We present two cases of patients diagnosed with a lung adenocarcinoma harboring rare EGFR mutations that received treatment with gefitinib. Mutational analysis was performed by polymerase chain reaction amplification of exons 18, 19, 20 and 21 of the EGFR gene and exon 1 of the KRAS gene followed by direct sequencing using BigDye 3.1 (Applied Biosystems, Foster City, CA) and analysis in a Genetic Analyzer 3500Dx (Applied Biosystems).

CASE 1

A white, 81-year-old, male exsmoker was diagnosed in July 2010 of a stage IIIA (T4N0M0) lung adenocarcinoma. Molecular analysis demonstrated the presence of a p.R836C mutation in exon 21 of the EGFR gene. He began first-line chemotherapy with a platinum doublet. Response assessment by computed tomography scan showed stabilization. However, because of symptomatic

progression he started treatment with gefitinib 250mg/day. In January 2011 he was admitted in hospital because of increasing dyspnoea and pain. The computed tomography scan showed progression of the tumor with appearance of pleural effusion. The study of the fluid confirmed the diagnosis of metastasis from adenocarcinoma harboring the p.R836C mutation on exon 21 (Fig. 1A). No p.T790M mutation on exon 20 or hepatocyte growth factor receptor (MET) amplification was detected. Progressive disease led to rapid deterioration of performance status and the patient died in early March 2011.

CASE 2

A white, 54-year-old, heavy smoker man was diagnosed in May 2011 with a lung adenocarcinoma stage IIIB (T4N2M0) not suitable for radical treatment. Evaluation of EGFR mutational status demonstrated the presence of the p.T854S mutation in exon 21. No wild-type allele was detected in this position, indicating that the mutation was either hemizygous or homozygous (Fig. 1B). He started treatment with gefitinib and early (6 weeks) assessment by computerized tomography demonstrated progressive disease. He received subsequent lines of treatment with stable disease as best response. Local and central nervous system progression led

to important clinical deterioration and death in March 2012.

DISCUSSION

The finding of *EGFR* mutations not previously described always confers the challenge of treatment decision. For this reason we report these cases of patients with infrequent *EGFR* mutations on exon 21 with rapid progression on gefitinib treatment. Upcoming results of newly available EGFR inhibitors such as afatinib or dacomitinib will show whether these drugs will be effective for this subset of patients.

Elisabet Mompradé, MD
Eduard Arriola, MD, PhD
Medical Oncology Department
Hospital del Mar
Barcelona, Spain

Luz Martínez-Avilés, MSc
Pathology Department
Hospital del Mar
Barcelona, Spain

REFERENCES

1. Lynch TJ, Bell DW, Sordella R, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness

- of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2004;350:2129–2139.
2. Mitsudomi T, Morita S, Yatabe Y, et al.; West Japan Oncology Group. Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. *Lancet Oncol* 2010;11:121–128.
3. Sequist LV, Waltman BA, Dias-Santagata D, et al. Genotypic and histological evolution of lung cancers acquiring resistance to EGFR inhibitors. *Sci Transl Med* 2011;3:75ra26.
4. Yatabe Y, Pao W, Jett JR. Encouragement to submit data of clinical response to EGFR-TKIs in patients with uncommon EGFR mutations. *J Thorac Oncol* 2012;7:775–776.